

The Impact of Outpatient Laboratory Alerting Mechanisms in Patients with Acute Kidney Injury

Nicole V. Tolan^{1,2†}, Salman Ahmed^{3†}, Tolumofe Terebo^{2,4}, Zain M. Virk², Athena K. Petrides^{1,2},
Jaime R. Ransohoff^{1,5}, Christiana A. Demetriou^{6,7}, Yvelynne P. Kelly⁸, Stacy E. F. Melanson^{1,2*},
Mallika L. Mendu^{2,4*}

†Equal contribution as first author, *Equal contribution as senior author

¹Department of Pathology, Brigham and Women's Hospital, Boston, MA, ²Harvard Medical School, Boston, MA, ³Section of General Internal Medicine, Department of Medicine, Baylor College of Medicine, Houston, TX, ⁴Department of Medicine, Renal Division, Brigham and Women's Hospital, Boston, MA, ⁵Department of Epidemiology, Bloomberg School of Public Health, Baltimore, MD, ⁶Department of Primary Care and Population Health, University of Nicosia Medical School, Nicosia, Cyprus, ⁷The Cyprus School of Molecular Medicine, The Cyprus Institute of Neurology and Genetics, Nicosia, Cyprus, ⁸Department of Critical Care Medicine, St. James Hospital, Trinity College, Dublin, Ireland.

Corresponding Author: Stacy E. F. Melanson, Brigham and Women's Hospital, 75 Francis Street, Amory 2, Boston, MA 02115, Tel: 617-732-6245, Email: semelanson@bwh.harvard.edu

Key Points

- An AKI alert attached to increasing creatinine results improved recognition of AKI and reduced the time to obtain a follow-up creatinine.
- The additive intervention of an email alert to the ordering provider increased the discontinuation of nephrotoxins.

Abstract

Background: Acute kidney injury (AKI) is an abrupt decrease in kidney function associated with significant morbidity and mortality. Electronic notifications of AKI have been utilized in hospitalized patients, but their efficacy in the outpatient setting is unclear.

Methods: We evaluated the impact of two outpatient interventions: an automated comment on increasing creatinine results (intervention I; 6 months; n=159) along with an email to the provider (intervention II; 3 months; n=105) compared to a control (baseline; 6 months; n=176). A comment was generated if a patient's creatinine increased by >0.5 mg/dL (previous creatinine ≤ 2.0 mg/dL) or by 50% (previous creatinine >2.0 mg/dL) within 180 days. Process measures included documentation of AKI and clinical actions. Clinical outcomes were defined as recovery from AKI within 7 days, prolonged AKI from 8 to 89 days, progression to CKD within 120 days.

Results: Providers were more likely to document AKI in interventions I ($p=0.004$; OR=2.80) and II ($p=0.01$; OR=2.66). Providers were also more likely to discontinue nephrotoxins in intervention II ($p<0.001$; OR= 4.88). The median time to follow-up creatinine trended shorter among patients with AKI documented (21 vs. 42 days; $p=0.11$). There were no significant differences in clinical outcomes.

Conclusions: An automated comment was associated with improved documented recognition of AKI and the additive intervention of an email alert was associated with increased discontinuation of nephrotoxins, but neither improved clinical outcomes. Translation of these findings into improved outcomes may require corresponding standardization of clinical practice protocols for managing AKI.

Introduction

Acute kidney injury (AKI) is an abrupt decrease in kidney function that may be due to a diverse array of etiologies.¹ The incidence of AKI varies from 22% to 57% depending on the care setting (e.g. outpatient, intensive care unit) and the criteria applied for case definition.^{2,3} AKI in the outpatient setting is common, occurring up to 31.5% of patients in one observational cohort.⁴ Possible factors associated with increasing prevalence of AKI include an aging population, rise in comorbidities that predispose to AKI (e.g. diabetes mellitus, hypertension, congestive heart failure, chronic kidney disease (CKD)), increased use of agents that may result in nephrotoxicity (e.g. chemotherapy, antimicrobials), and increasing frequency of invasive and surgical procedures.^{5,6} Each episode of AKI is consequential, and is associated with outcomes including mortality, cardiovascular events, CKD, and end stage renal disease (ESRD).^{7,8}

Identification of AKI allows clinicians to follow consensus guideline recommendations for the appropriate management of patients with AKI including assessment of medication dosing, avoiding nephrotoxic exposures, and managing fluid and electrolytes.⁸ However, there is considerable variability in clinician and healthcare systems adherence to these guidelines to prevent, diagnose, and treat AKI, which may account for the high rates of adverse outcomes.⁸

Electronic health records (EHR) can enable effective population surveillance across many fields of medicine. Related to AKI, EHR-based alert systems may allow timely recognition of AKI, guide appropriate clinical action, and improve efforts to prevent AKI.^{9,10} Most reports of electronic notification of AKI have focused on in-hospital AKI events.¹¹⁻¹³ However, early identification and management of AKI in the outpatient setting may prevent worsening AKI and the associated consequences of severe and/or prolonged AKI. Kidney function in the 72-hour period immediately following AKI in hospitalized patients is associated with specific clinical

outcomes: the development or progression of CKD, initiation of long-term dialysis, or death from any cause within a median follow-up of approximately 4.7 years.¹⁴ In this study, we implemented sequential AKI alert interventions for outpatient providers and determined if the interventions were associated with increased documentation of AKI and related clinical actions that could improve clinical outcomes.

Materials and Methods

Study Setting

Brigham and Women's Hospital is a 793-bed tertiary care hospital located in Boston, MA. The clinical laboratories process approximately 3 million specimens/year and perform approximately 275 outpatient plasma creatinine tests per day; with a baseline rate of outpatient AKI of 1-2%. Estimated glomerular filtration rate (eGFR) has been calculated using the CKD-EPI equation since March 6, 2018. Prior to that, the MDRD equation was utilized. The laboratory information system (LIS) is Sunquest (SQ; Sunquest Information Systems, Inc., Tucson, AZ) and electronic health record (EHR) is Epic (Epic Healthcare Systems, Verona, WI). The project was conducted under Mass General Brigham Institutional Review Board exemption as meeting the requirements of quality improvement research.

Sample Selection, Interventions and Exclusion Criteria

The study was divided into three intervals (Figure 1): a baseline or control population without any AKI alert (a 6-month period from 2/26/2018 to 8/24/2018), an automated result comment attached to increasing creatinine results (intervention I; a 6-month period from 9/17/2018 to 3/15/2019) and a manual email to the ordering provider regarding the result comment in addition to the automated AKI comment (intervention II; a 3-month period from 7/8/2019 to 9/7/2019). Intervention II was limited to 3 months because of the effort required by the laboratory directors to perform daily case review and statistical power had been achieved in this timeframe.

Utilizing a calculation built in our LIS, an automated creatinine result comment was generated if a patient's creatinine increased by >0.5 mg/dL if the previous creatinine was ≤ 2.0

mg/dL, or increased by 50% if the previous creatinine was >2.0 mg/dL (Figure 2a). The most recent creatinine within the past 180 days was used as the baseline creatinine from which to calculate the delta. In both intervention I and II, result comments (Figure 2b) were appended to creatinine results for outpatients when these criteria were met. The result comment was not applied to patient results from dialysis locations or if no creatinine was available in the prior 180 days. In intervention II, an automated crystal report was built to identify these cases of potential AKI and was generated three times per day at approximately 2am, 10am and 6pm. One of two laboratory directors (N.T., S.M.) manually reviewed each case and emailed the ordering provider within 24 hours of the automated result comment to alert them of the potential risk of AKI and included information regarding the previous and current creatinine results (Figure 2c).

Prior to analysis, a manual review was conducted by one of two laboratorians (N.T, S.M.) to exclude cases from each interval if 1) the patient had the diagnosis of ESRD or was on renal replacement therapy (RRT), 2) the patient had a creatinine resulted at an outside hospital (OSH) between the flagged and previous creatinine at BWH and the patient no longer met the criteria for AKI (Figure 2a), or 3) the patient was hospitalized for reasons other than AKI the day the flagged creatinine was resulted (Figure 1). If the patient had more than one flagged creatinine within a 2-week period, only the earliest flagged creatinine was included. In intervention II, the laboratory director did not email the provider if there was a note in the chart that the provider was already aware of the results at the time of review.

Process Measures

Process measures included documentation of AKI and clinical action taken. Clinical actions were only analyzed if the provider documented AKI. We also determined if the time to obtain a follow-up creatinine differed between providers that did and did not document AKI.

Furthermore, we investigated whether documentation and clinical actions differed between ordering provider services and whether the interventions changed provider behavior related to these process measures.

Outcome Measures

Given the outpatient population studied, we defined our clinical outcomes as 1) recovery from AKI within 7 days, 2) prolonged AKI between 8 to 89 days and 3) progression to CKD from 90 to 120 days after the AKI alert. Recovery was defined as a decrease in creatinine of ≥ 0.3 mg/dL or 25% from peak creatinine. Prolonged AKI was defined as a creatinine that did not meet the recovery criteria. Progression to CKD was defined as two eGFR < 60 mL/min/1.73m² separated by 90 to 120 days.

Medical Record Review for Process Measures and Clinical Outcomes

Medical record review to obtain process measures of AKI and clinical actions was completed independently by one of two clinicians (S.A., Z.V.) using a standardized review instrument. To assess reliability of the medical record review, 10% of cases were reviewed by the other clinician using the same standardized template and analyzed for concordance.

The following parameters were obtained during medical record review (Supplemental Table 1): documented recognition of AKI and date, documented clinical action and date, type of clinical action (i.e., repeat labs, encourage hydration, discontinuation of nephrotoxins, change dosing, nephrology consult, hospitalization), presence of underlying CKD and cause of AKI. Each parameter, except dates and cause of AKI, was captured as binary data, having occurred or not. Only documentation and clinical action occurring within the two weeks following the AKI alert was included.

AKI cause was further categorized as hypovolemia (including overdiuresis), cardiorenal syndrome, drugs (including contrast) or other, to allow for binary analysis. Patients with more than one documented AKI cause were also categorized as multifactorial. If a patient was hospitalized, the date of admission and length of stay were recorded. The time (in days) between the creatinine result comment and the documentation of each recognition of AKI and clinical action, was determined.

LIS and EHR Data

Patient demographics (i.e. age, gender, race) and follow-up creatinine measurement(s) from 0 to 120 days post AKI alert were obtained from our LIS. The time to first follow-up creatinine was also calculated. The ordering provider and their clinical service was obtained electronically from the specialty description field in our EHR. Provider services were grouped into five categories: internal medicine (study total n= 187; average across intervals n=62), cardiology (total n=83; average n=28), pulmonary (total n =71; average n=24), nephrology (total n=50; average n=17), and surgical services/transplant (total n=49; average n=16), Dermatology, endocrinology, gastroenterology, infectious disease, primary care and family medicine were grouped with internal medicine and anesthesiology, urology, oncology, cardiac transplant and pulmonary transplant were considered surgical services/transplant.

Statistical Methods

Patient demographics and characteristics were compared between intervention categories using the Pearson's chi-squared (χ^2) test for categorical outcomes, the Fisher's exact test for categorical outcomes when cell counts were <5, and the Wilcoxon rank-sum test for non-normally distributed continuous variables. A Cohen's kappa statistical test was performed to

assess agreement between two reviewers for all process measures and AKI causes. Logistic regression adjusted for age, sex, race, flagged creatinine, and AKI causes (drugs including contrast, multifactorial and no documented cause) was used to assess differences in the likelihood of each process measure between intervention categories. P-values are reported both for the comparison of each intervention category to the baseline, as well as for the trend. Based on the sample size of this investigation, the minimum % increase in the dichotomous outcomes that could be detected as statistically significant – in any comparison – was 15%, assuming a power of 0.80 and an alpha level of 5%. All analyses were performed using SAS version 9.4 (SAS Institute, Cary, NC) & STATA version SE15 (StataCorp. 2015. *Stata Statistical Software: Release 14*. College Station, TX: StataCorp LP).

Results

Study Characteristics and Patient Demographics

The overall dataset curation, including the number of excluded cases and reasons for exclusions, are shown in Figure 1. There was no difference in excluded cases in each interval ($p = 0.29$). The majority of creatinine result comments were based on calculation #1 (Figure 2a) with only 2.8%, 0.7% and 3.8% flagging based on calculation #2 in baseline, intervention I and intervention II; respectively. The rates of recurrent AKI were low and similar in each interval ranging from 2 to 7% ($p = 0.14$). Table 1 lists the patient demographics and characteristics in each interval of the study. There was a difference in Hispanic race ($p = 0.01$), median creatinine results with automated comments (i.e. median flagged creatinine) ($p = 0.02$), and the AKI causes of drugs including contrast ($p < 0.001$), multifactorial ($p < 0.001$), and no documented cause ($p = 0.001$) between the intervals; requiring normalization in subsequent analyses.

Process Measures

Providers were more likely to document AKI in both intervention I ($p = 0.004$; OR = 2.80) and intervention II ($p = 0.01$; OR = 2.66) compared to baseline (Table 2 and 3). The trend p -value for documentation of AKI was also significant ($p = 0.006$; OR = 1.74). Providers were also more likely to discontinue nephrotoxins in intervention II compared to baseline ($p < 0.001$; OR = 4.88), and there was a significant trend across intervals ($p < 0.001$; OR = 2.22) (Table 3). However, there was no significant difference in the type of nephrotoxins discontinued across the intervals (Table 4). The AKI alert interventions did not significantly impact any other clinical action (repeat labs, encourage hydration, changing dosing, nephrology consult or hospitalization) (Table 2 and 3).

Although not statistically significant, the median time to obtain a follow-up creatinine across all intervals by providers who documented AKI and did not document AKI was 21 days (IQR 68) and 42 days (IQR 63); respectively ($p = 0.11$).

Clinical Outcomes

The AKI alert interventions did not significantly impact the clinical outcomes of AKI recovery, prolonged AKI, or progression to CKD when compared to baseline (Table 3). The number of follow up creatinine results at 0 to 7 days, 8 to 89 days and 90 to 120 days was similar across the interventions ($p = 0.07$, $p = 0.13$, $p = 0.07$; respectively) (Table 2). Similarly, the presence of underlying CKD in patients with CKD progression was 87% in baseline, 84% in intervention I and 85% in intervention II ($p = 0.91$).

Difference Among Clinical Services

Regardless of the interval, surgical services/transplant were less likely than any other service to document both AKI ($p < 0.001$; OR = 0.26, 95% CI (0.12, 0.57)) and clinical action ($p < 0.001$; OR = 0.20, 95% CI (0.11, 0.37)). In contrast, the pulmonary service was more likely to both document AKI ($p = 0.01$; OR = 3.18, 95% CI (1.29, 7.85)) and clinical action ($p = 0.03$, OR = 2.12, 95% CI (1.09, 4.09)).

Internal medicine providers were more likely to document AKI in both intervention I ($p = 0.02$, OR = 3.97, 95% CI (1.27, 11.85)) and intervention II ($p = 0.008$, OR = 6.07, 95% CI (1.61, 22.90)) compared to baseline with a trend p -value = 0.004 (Table 5). There were no significant changes in documentation of AKI in other services. Internal medicine providers were also more likely to document clinical action. There was difference between baseline and intervention II ($p = 0.02$, OR = 5.76, 95% CI (1.30, 25.69)) as well as a significant trend ($p =$

0.02, OR = 2.41, 95% CI (1.15, 5.03)) (Table 5). The interventions did not significantly impact clinical action in the other services.

Reviewer Agreement

Reviewers achieved almost perfect agreement for hospitalization (K = 0.811) and cardiorenal syndrome (K = 0.847). Reviewers achieved substantial agreement on documented recognition of AKI (K = 0.627), encourage hydration (K = 0.677), discontinuation of nephrotoxins (K = 0.738), hypovolemia (K = 0.761) and no documented cause of AKI (K = 0.610). Moderate agreement was achieved for documented clinical action (K = 0.502) and other as cause for AKI (K = 0.453). Reviewers achieved fair agreement for repeat labs (K = 0.399), change dosing (K = 0.381), nephrology consult (K = 0.370) and drugs (including contrast) (K = 0.361). All components of the medical record review scored a K-value ≥ 0.36 corresponding to fair or better and $\geq 75\%$ agreement.

Discussion

In this intervention-based cohort study, we found that an EHR-based automated comment displaying with increasing creatinine results was associated with increased documented recognition of AKI, particularly by internal medicine providers, and a decreased time to obtain follow up creatinine measurements. The additive intervention of an email notification sent by a laboratory director to the ordering provider was associated with an increase in the discontinuation of nephrotoxins. To our knowledge, this is the first study to examine the impact of AKI alerts and define clinical outcomes specific for the outpatient setting.

Previous studies on the impact of AKI alerts were performed on hospitalized patients and reported variable impacts on clinical outcomes.^{8-13,15-17} Aiyegbushi et al. performed a study on hospitalized patients and showed that an AKI alert was associated with a dramatic reduction in the repeat testing interval for creatinine.¹⁵ However, Wilson et al. in a single-center, randomized controlled trial of more than 23,000 hospitalized adult patients found that an AKI alert had no impact on the clinical outcomes of peak creatinine concentration, initiation of RRT, or death during the current hospital admission or within 30 days.¹¹ In that study, the intervention group received a text page alert indicating potential AKI, following the 2012 The Kidney Disease: Improving Global Outcomes (KDIGO) consensus definition for AKI.⁸ Another randomized clinical trial also showed no impact of a pop-up alert for AKI in hospitalized patients on clinical outcomes.¹³

Laboratories usually have well-defined processes for communicating immediately life-threatening or critical results such as hyperkalemia, but processes and algorithms for communicating significant changes in results such as creatinine are not routinely implemented or standardized across institutions. The effectiveness of electronic notifications is variable and

reportedly more successful if alerts are simplistic, evidence-based and only triggered at the appropriate time.^{18,19} An effective AKI alert should include both an accurate baseline creatinine, the definition of which is a subject of debate,^{8,20-22} and a change in creatinine that is sensitive, yet specific for AKI.⁸ According to Siew et al. the mean outpatient creatinine from the past 7-365 days is the best approximation of the baseline creatinine concentration.²³ However, not all laboratories have the time, resources, and technologic infrastructure to implement AKI alerts using running mean creatinine values. The 2012 KDIGO consensus definition for AKI is described as a change of ≥ 0.3 mg/dL in the prior 48 hours,⁸ but a high number of false-positive AKI alerts have been seen using these guidelines, particularly in patients with underlying CKD and higher baseline creatinine concentrations.^{24,25}

In our study, providers were significantly more likely to document recognition of AKI in both interventions. The time to obtain a follow-up creatinine was 21 days if AKI was documented as compared to 42 days if AKI was not documented across all interventions. Although not statistically significant, the difference is clinically significant and similar to the findings of Aiyegbushi et al.¹⁵ It is notable that more participants had a follow-up creatinine at 8-89 days as opposed to the other time windows which may have implications on recovery from AKI. However, we hypothesize that this pattern reflects outpatient AKI management involving a period of observation after a clinical action. Providers were also significantly more likely to discontinue nephrotoxins in intervention II compared to baseline. Although pre-renal causes of AKI including hypovolemia are common, as seen in our study, the impact of discontinuation of nephrotoxins can be significant. A larger sample size was likely necessary to show an impact of discontinuing nephrotoxins on recovery from AKI, prolonged AKI and progression to CKD. The impact of interventions on nephrology consults could not be assessed as there were no consults

in intervention II. This may suggest that the email increased the confidence of non-nephrology providers in diagnosing and managing AKI. Overall, our findings suggest that a result comment alone improved documented recognition of AKI and consequentially decreased the time to obtain a follow up creatinine. However, an email was necessary to increase the discontinuation of nephrotoxins and possibly reduce nephrology consults.

Across all intervals, surgical and transplant services were least likely to document AKI and clinical actions, while pulmonary was most likely to document AKI and clinical actions. This may be due to the availability of standardized protocols for documentation that have been implemented by the pulmonary team at our institution. Internal medicine providers, the largest group of providers in our study, were significantly more likely to document recognition of AKI in both intervention I and II and more likely to document clinical action in intervention II. This suggests that internal medicine, unlike other services, may be the most responsive to alerts and managing AKI in the outpatient setting, and could reflect greater clinical expertise and education about AKI as opposed to other provider groups. Furthermore, it may be more effective to alert the primary care provider of AKI in addition to or instead of the ordering provider regardless of service.

We speculate that we did not see an impact of our interventions on clinical outcomes for several reasons. First, we had a limited sample size, constrained based on the number of patients with outpatient creatinine obtained during the time periods outlined, which likely limited our power to observe differences in clinical outcomes. Second, we were limited to data within our EHR, which means that patients may have had labs checked indicating recovery or CKD development that we were unable to capture. Third, we did not capture data on the patient population, medical and social complexity, which can impact the ability to mitigate the

consequences of AKI. Unlike previously published studies on AKI alerts, our patients had AKI in the outpatient setting, had higher baseline kidney function and were less likely to be hospitalized as a result of AKI (<15% in each interval; Table 2). Therefore, providers may have been focused on managing other co-morbidities. Fourth, we did not educate providers about strategies to manage AKI and about the alert interventions themselves. We recognize that AKI alerts alone may not be enough to improve clinical outcomes. While the impact of our interventions on process measures including AKI documentation and discontinuation of nephrotoxins was promising, integrated education, such as standardized clinical practice guidelines, daily review of alerts by a nephrologist, and/or more robust alerts such as a text page may be necessary to affect process measures and clinical outcomes, particularly for providers outside of internal medicine.

Our study has several limitations. First, the study was performed at a large academic medical center and results may not be generalizable. Second, our study was not randomized and interventions were sequential, which limits causality, though there were no other efforts undertaken at the same time to improve AKI. Utilizing the available functionality in our LIS, we evaluated each creatinine against the most recent creatinine measurement in the previous 180 days and our criteria for AKI were modified for our outpatient population. Therefore, the determination of baseline was not optimal, leading to both false-positive and negative AKI alerts.²⁵ However, feedback from providers receiving the alerts was generally positive (e.g. “This new notification system is great and will improve patient safety”). Furthermore, no concerns were raised about the accuracy or timing of the alert. Lastly, in intervention II we did not email the ordering provider with the AKI alert if the provider noted in the EHR that they were aware of

the increasing creatinine. Therefore, we may have underestimated the impact of intervention II throughout our analyses.

Based on our findings the laboratory continues to append an automated comment to increasing creatinine results. We encourage nephrologists to collaborate with their laboratory colleagues and implement a calculation in their LIS that appends a comment to increasing creatinine results indicative of AKI in the outpatient setting. Additional studies are required to confirm the optimal baseline creatinine, the thresholds for AKI alert in the outpatient population, the most effective mechanism of notification (such as a text page) and which providers to send the notification. The utility of alerts to trigger a review of nephrotoxins and prevent AKI should also be explored. Finally, developing education that accompanies the alert and consensus guidelines for management of AKI for outpatients will be important.

Disclosures: N. Tolan reports the following: Research Funding: Abbott Diabetes Care, Biomerieux; Scientific Advisor or Membership: Journal of Applied Laboratory Medicine, College of American Pathologists, American Association for Clinical Chemistry. S. Ahmed reports the following: Ownership Interest: The Kidney Health and Preventive Medicine Institute. M. Mendu reports the following: Consultancy Agreements: Bayer AG. The remaining authors have nothing to disclose.

Funding: None

Author Contributions: N Tolan: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Supervision; Validation; Visualization; Writing - original draft; Writing - review and editing

S Ahmed: Data curation; Formal analysis; Investigation; Validation; Visualization; Writing - review and editing

T Terebo: Data curation; Project administration; Validation; Visualization; Writing - review and editing

Z Virk: Data curation; Investigation; Validation; Visualization; Writing - review and editing

A Petrides: Conceptualization; Formal analysis; Investigation; Methodology; Writing - review and editing

J Ransohoff: Data curation; Project administration; Validation; Visualization; Writing - review and editing

C Demetriou: Formal analysis; Investigation; Methodology; Validation; Writing - review and editing

Y Kelly: Conceptualization; Data curation; Methodology; Supervision; Validation; Writing - review and editing

S Melanson: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Resources; Supervision; Validation; Visualization; Writing - original draft; Writing - review and editing

M Mendu: Conceptualization; Investigation; Methodology; Supervision; Validation; Visualization; Writing - review and editing

Supplemental Material Table of Contents

Supplemental Table 1. Definitions of Parameters Collected During Medical Record Review

References

1. Hoste EAJ, Kellum JA, Selby NM, Zarbock A, Palevsky PM, Bagshaw SM, Goldstein SL, Cerdá J, Chawla LS: Global epidemiology and outcomes of acute kidney injury. *Nat Rev Nephrol* 14: 607–625, 2018
2. Susantitaphong P, Cruz DN, Cerda J, Abulfaraj M, Alqahtani F, Koulouridis I, Jaber BL: World Incidence of AKI: A Meta-Analysis. *Clin J Am Soc Nephrol* 8: 1482–1493, 2013
3. Hoste EAJ, Bagshaw SM, Bellomo R, Cely CM, Colman R, Cruz DN, Edipidis K, Forni LG, Gomersall CD, Govil D, Honoré PM, Joannes-Boyau O, Joannidis M, Korhonen A-M, Lavrentieva A, Mehta RL, Palevsky P, Roessler E, Ronco C, Uchino S, Vazquez JA, Vidal Andrade E, Webb S, Kellum JA: Epidemiology of acute kidney injury in critically ill patients: the multinational AKI-EPI study. *Intensive Care Med* 41: 1411–1423, 2015
4. Yeh H-C, Ting I-W, Huang H-C, Chiang H-Y, Kuo C-C: Acute Kidney Injury in the Outpatient Setting Associates with Risk of End-Stage Renal Disease and Death in Patients with CKD. *Sci Rep* 9: 17658, 2019
5. Siew ED, Davenport A: The growth of acute kidney injury: a rising tide or just closer attention to detail? *Kidney Int* 87: 46–61, 2015
6. Waikar SS, Liu KD, Chertow GM: The incidence and prognostic significance of acute kidney injury. *Curr Opin Nephrol Hypertens* 16: 227–236, 2007
7. Xue JL, Daniels F, Star RA, Kimmel PL, Eggers PW, Molitoris BA, Himmelfarb J, Collins AJ: Incidence and Mortality of Acute Renal Failure in Medicare Beneficiaries, 1992 to 2001. *J Am Soc Nephrol* 17: 1135–1142, 2006
8. Khwaja A: KDIGO Clinical Practice Guidelines for Acute Kidney Injury. *Nephron Clin Pract* 120: c179–c184, 2012
9. James MT, Hobson CE, Darmon M, Mohan S, Hudson D, Goldstein SL, Ronco C, Kellum JA, Bagshaw SM, For the Acute Dialysis Quality Initiative (ADQI) Consensus Group: Applications for Detection of Acute Kidney Injury Using Electronic Medical Records and Clinical Information Systems: Workgroup Statements from the 15th ADQI Consensus Conference. *Can J Kidney Health Dis* 3: 100, 2016
10. Hoste EAJ, Kashani K, Gibney N, Wilson FP, Ronco C, Goldstein SL, Kellum JA, Bagshaw SM: Impact of Electronic-Alerting of Acute Kidney Injury: Workgroup Statements from the 15th ADQI Consensus Conference. *Can J Kidney Health Dis* 3: 101, 2016
11. Wilson FP, Shashaty M, Testani J, Aqeel I, Borovskiy Y, Ellenberg SS, Feldman HI, Fernandez H, Gitelman Y, Lin J, Negoianu D, Parikh CR, Reese PP, Urbani R, Fuchs B: Automated, electronic alerts for acute kidney injury: a single-blind, parallel-group, randomised controlled trial. *Lancet* 385: 1966–1974, 2015

12. Lachance P, Villeneuve P-M, Rewa OG, Wilson FP, Selby NM, Featherstone RM, Bagshaw SM: Association between e-alert implementation for detection of acute kidney injury and outcomes: a systematic review. *Nephrol Dial Transplant* 32: 265-272, 2017
13. Wilson FP, Martin M, Yamamoto Y, Partridge C, Moreira E, Arora T, Biswas A, Feldman H, Garg AX, Greenberg JH, Hinchcliff M, Latham S, Li F, Lin H, Mansour SG, Moledina DG, Palevsky PM, Parikh CR, Simonov M, Testani J, Ugwuowo U: Electronic health record alerts for acute kidney injury: multicenter, randomized clinical trial. *BMJ* 372: m4786, 2021
14. Bhatraju PK, Zelnick LR, Chinchilli VM, Moledina DG, Coca SG, Parikh CR, Garg AX, Hsu C, Go AS, Liu KD, Ikizler TA, Siew ED, Kaufman JS, Kimmel PL, Himmelfarb J, Wurfel MM: Association Between Early Recovery of Kidney Function After Acute Kidney Injury and Long-term Clinical Outcomes. *JAMA Netw Open* 3: e202682, 2020
15. Aiyegbusi O, Witham MD, Lim M, Gauld G, Bell S: Impact of introducing electronic acute kidney injury alerts in primary care. *Clin Kidney J* 12: 253–257, 2019
16. Predecki M, Blacker E, Sadeghi-Alavijeh O, Edwards R, Montgomery H, Gillis S, Harber M: Improving outcomes in patients with Acute Kidney Injury: the impact of hospital based automated AKI alerts. *Postgrad Med J* 92: 9–13, 2016
17. Sykes L, Nipah R, Kalra P, Green D: A narrative review of the impact of interventions in acute kidney injury. *J Nephrol* 31: 523–535, 2018
18. Thomas ME, Blaine C, Dawnay A, Devonald MAJ, Ftouh S, Laing C, Latchem S, Lewington A, Milford DV, Ostermann M: The definition of acute kidney injury and its use in practice. *Kidney Int* 87: 62–73, 2015
19. Bellomo R, Ronco C, Kellum JA, Mehta RL, Palevsky P, the ADQI workgroup: Acute renal failure – definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care* 8: R204, 2004
20. Mehta RL, Kellum JA, Shah SV, Molitoris BA, Ronco C, Warnock DG, Levin A, the Acute Kidney Injury Network: Acute Kidney Injury Network: report of an initiative to improve outcomes in acute kidney injury. *Crit Care* 11: R31, 2007
21. Embi PJ, Leonard AC: Evaluating alert fatigue over time to EHR-based clinical trial alerts: findings from a randomized controlled study. *J Am Med Inform Assoc* 19: e145-148, 2012
22. Bates DW, Kuperman GJ, Wang S, Gandhi T, Kittler A, Volk L, Spurr C, Khorasani R, Tanasijevic M, Middleton B: Ten Commandments for Effective Clinical Decision Support: Making the Practice of Evidence-based Medicine a Reality. *J Am Med Inform Assoc* 10: 523–530, 2003
23. Siew ED, Ikizler TA, Matheny ME, Shi Y, Schildcrout JS, Danciu I, Dwyer JP, Srichai M, Hung AM, Smith JP, Peterson JF: Estimating Baseline Kidney Function in Hospitalized Patients with Impaired Kidney Function. *Clin J Am Soc Nephrol* 7: 712–719, 2012

24. Lin J, Fernandez H, Shashaty MGS, Negoianu D, Testani JM, Berns JS, Parikh CR, Wilson FP: False-Positive Rate of AKI Using Consensus Creatinine-Based Criteria. *Clin J Am Soc Nephrol* 10: 1723–1731, 2015
25. Joe M E-K, Melanie P H, Graham RD J, Edmund J L, Chirag R P, et al.: AACC Guidance Document on Laboratory Investigation of Acute Kidney Injury. *J Appl Lab Med* *in press*, 2021

Table 1. Patient Demographics and Characteristics

	Baseline (n=176)	Intervention I (n=159)	Intervention II (n=105)	p-Value
Male, % (n)	55.1% (97)	49.1% (78)	53.3% (56)	0.53
Age, Mean ± SD	65 ± 14.0	65 ± 13.3	66 ± 15.9	0.27
White, % (n)	75.6% (133)	66.0% (105)	72.4% (76)	0.15
Black, % (n)	11.4% (20)	18.2% (29)	18.1% (19)	0.15
Hispanic, % (n)	11.4% (20)	5.0% (8)	2.9% (3)	0.01
Asian, % (n)	1.7% (3)	1.3% (2)	1.0% (1)	0.86
Flagged Cr, Median, (IQR)	2.13 (0.68)	1.94 (0.63)	2.08 (0.67)	0.02
Underlying CKD % (n)	67.6% (119)	67.3% (107)	65.7% (69)	0.94
Hypovolemia, % (n)	53.4% (94)	50% (80)	51.4% (54)	0.92
Cardiorenal Syndrome, % (n)	8.5% (15)	2.5% (4)	7.6% (8)	0.06
Drugs (Incl. Contrast), % (n)	30.1% (53)	15.6% (25)	8.6% (9)	p < 0.001
Multifactorial, % (n)	18.8% (33)	18.1% (13)	3.8% (4)	p < 0.001
No Documented Cause, % (n)	14.2% (25)	30.8% (49)	25.7% (27)	0.001
Other, % (n)	11.9% (21)	8.1% (13)	10.5% (11)	0.52

Cr = creatinine, CKD = chronic kidney disease, Incl. = Including

Table 2. Clinical Actions Taken During Each Interval

Clinical Action	Baseline (n=123)	Intervention I (n=109)	Intervention II (n=76)
Repeat Labs, % (n)	80.5% (99)	82.6% (90)	82.9% (63)
Follow up Cr at 0-7 days, % (n)	34.7% (61)	41.5% (66)	27.6% (29)
Follow up Cr at 8-89 days, % (n)	79.0% (139)	71.1% (113)	69.5% (73)
Follow up Cr at 90-120 days, % (n)	29.5% (87)	40.9% (65)	36.2% (38)
Encourage Hydration, % (n)	41.5% (51)	47.7% (52)	48.7% (37)
Discontinuation of Nephrotoxins, % (n)	17.9% (22)	19.3% (21)	38.2% (29)
Change Dosing, % (n)	35.0% (43)	35.8% (39)	25.0% (19)
Nephrology Consult, % (n)	7.3% (9)	1.8% (2)	0% (0)
Hospitalization, % (n)	14.6% (18)	11.9% (13)	14.5% (11)

Cr = creatinine

Table 3. Process Measures and Clinical Outcomes

	Intervention I (n=159)	Intervention II (n=105)	Trend p-value
Documented AKI, n, OR (95%CI)	1112.80(1.38, 5.67)	772.66(1.21, 5.81)	0.006
Clinical Action, OR(95%CI)	2.54(0.50, 12.83)	2.01(0.35, 11.73)	0.37
Repeat Labs, OR(95%CI)	1.60(0.77, 3.30)	1.69(0.77, 3.73)	0.17
Encourage Hydration, OR(95%CI)	1.57(0.87, 2.83)	1.82(0.95, 3.48)	0.06
Discontinuation of Nephrotoxins, p value; OR(95%CI)	1.60(0.76, 3.35)	4.88(2.27, 10.50)	<0.001
Change Dosing, OR(95%CI)	1.02(0.57, 1.81)	0.60(0.31, 1.17)	0.17
Nephrology Consult, OR(95%CI)	0.33(0.07, 1.70)	--	0.05
Hospitalization, OR(95%CI)	0.95(0.42, 2.17)	1.20(0.50, 2.90)	0.71
Recovery within 7 days, n, OR(95%CI)	34 0.60(0.29, 1.2)	18 0.92(0.37, 2.30)	0.61
Prolonged AKI, n, OR(95%CI)	33 0.92(0.54, 1.58)	19 0.79(0.42, 1.48)	0.46
Progression to CKD, n, OR(95%CI)	49 0.91(0.43, 1.94)	33 1.97(0.680, 5.72)	0.31

AKI = acute kidney injury, OR = odds ratio, CI = confidence interval

Table 4. Discontinuation of Nephrotoxins in Each Interval

Class of Nephrotoxin Discontinued	Baseline (n=22)	Intervention I (n=21)	Intervention II (n=29)
Diuretic, % (n)	50% (11)	52% (11)	72% (21)
ACE inhibitor/ARB, % (n)	32% (7)	19% (4)	24% (7)
Immunosuppressant, % (n)	18% (4)	5% (1)	7% (2)
Antibiotics, % (n)	14% (3)	10% (2)	0% (0)
NSAID, % (n)	9% (2)	19% (4)	3% (1)
Anti-Diabetic, % (n)	0% (0)	10% (2)	7% (2)
Anti-Gout, % (n)	0% (0)	10% (2)	0% (0)
Anti-Hypertensive, % (n)	0% (0)	5% (1)	3% (1)
Anti-Arrhythmic, % (n)	0% (0)	5% (1)	0% (0)

ACE = angiotensin converting enzyme, ARB = angiotensin II receptor blocker; NSAID = non-steroidal anti-inflammatory

Table 5. Documentation of Acute Kidney Injury and Clinical Action by Clinical Service

	Intervention I (n=159)	Intervention II (n=105)	Trend p-value
Documented Recognition of AKI			
Internal Medicine (n=189) OR(95% CI)	3.90(1.27, 11.85)	6.07(1.61, 22.90)	0.004
Cardiology (n=83) OR(95% CI)	2.35(0.43, 12.75)	1.11(0.16, 7.83)	0.72
Nephrology (n=49) OR(95% CI)	1.52(0.15, 15.28)	0.07(0.002,2.045)	0.13
Documented Clinical Action			
Internal Medicine (n=189) p value*, OR(95% CI)	2.42(0.72, 8.13)	5.76(1.30, 25.69)	0.02
Cardiology (n=83) OR(95% CI)	2.02(0.38, 10.83)	1.00(0.16, 6.02)	0.91
Nephrology (n=49) OR(95% CI)	2.91(.035, 23.91)	0.52(0.49, 5.56)	0.74

AKI = acute kidney injury, OR = odds ratio, CI = confidence interval

Surgical and pulmonary services are not shown due to low numbers.

Figure Legends

Figure 1. Study Design. The three intervals of the study, associated interventions, number of and reason for exclusion and final case count are depicted.

AKI = acute kidney injury, Cr = creatinine, OSH = outside hospital, ESRD = end stage renal disease, RRT = renal replacement therapy, CKD = chronic kidney disease

*Patients may have more than one reason for exclusion

Figure 2. Study Interventions. a) The LIS calculations to flag outpatient creatinine in patients with possible acute kidney injury are shown. b) The comment that displayed in our EHR in intervention I and intervention II is depicted. c) The email template sent to the ordering provider in intervention II is shown.

Figure 1




Figure 1

Figure 2

a.

Calculation #1	Previous creatinine \leq 2.0 mg/dL AND Increase of $>$ 0.5 mg/dl within 180 days
Calculation #2	Previous creatinine $>$ 2.0 mg/dL AND Increase of 50% within 180 days

b.

Basic metabolic panel (BWH ,BWF ,DFCI ,MGH ,NWH ,MEEI ,NSMC ,SRH) Order: 576939233
 Collected: 8/23/2019 14:58 Status: Final result Visible to patient: No (not released) Next appt: None
 Specimen Information: Blood

ⓘ Newer results are available. Click to view them now.

	Ref Range & Units	8/23/19 1458	8/23/19 1328	8/14/19 1109	5/23/19 1411
SODIUM	136 - 145 mmol/L	141		142	140
POTASSIUM	3.4 - 5.1 mmol/L	5.0		4.0	4.5
CHLORIDE	98 - 107 mmol/L	106		105	103
CO2	22 - 31 mmol/L	20 ▼		26	27
BUN	6 - 23 mg/dL	18		13	19
CREATININE	0.50 - 1.20 mg/dL	1.39 ▲		0.87	0.89

Comment: **INCREASING CREATININE, POSSIBLE ACUTE KIDNEY INJURY**

- c. Your patient (last name, first name, MRN XXXX) has **INCREASING CREATININE AND POSSIBLE ACUTE KIDNEY INJURY** based on hospital guidelines:
 Previous creatinine (insert date): XXXX mg/dL
 Most recent creatinine (insert date): XXXX mg/dL

Figure 2

Supplemental Table of Contents

Supplemental Table 1. Definitions of Parameters Collected During Medical Record Review

Supplemental Table 1. Definitions of Parameters Collected During Medical Record Review

Parameter	Definition*
Documented AKI	Any clinician acknowledged AKI in the EHR in a visit note (any visit note, not limited to nephrology) or patient communication within two weeks of the AKI alert.
Documented Clinical Action	Any clinician documented one or more of the clinical actions below in the EHR in a visit note (any visit note, not limited to nephrology) or patient communication within two weeks of the AKI alert.
Repeat Labs	Any clinician noted that laboratory testing should be repeated.
Encourage Hydration	Any clinician noted that the patient should hydrate, increase fluid intake or present to a care facility for intravenous hydration.
Discontinue Medications	Any clinician noted that the patient should stop taking a medication.
Change Dosing	Any clinician noted that the patient should change the dosing of a medication or stop taking a medication temporarily. The temporary hold was included in “change dosing” because the authors intended for the “discontinue medications” field to reflect permanent or indefinite discontinuation of a medication.
Nephrology Consult	A nephrology consult occurred (as evidenced by a nephrology visit note) within 2 weeks of the AKI alert. Alternatively, there was documentation that the patient had a nephrology consult at an outside facility within 2 weeks of the AKI alert.
Hospitalization	The patient was hospitalized specifically for evaluation and management of AKI. The following criteria had to be met: 1) patient was admitted to observation or inpatient status at a Mass General Brigham inpatient facility and 2) the admission occurred after the AKI alert. At least one instance of documentation from that admission, e.g. emergency department expect note or inpatient history and physical, listed AKI as the primary problem, chief complaint, or reason for hospitalization.
Underlying CKD	The clinician noted in the EHR that the patient had a diagnosis of CKD in a visit note, patient communication, or problem list prior to the AKI alert. Alternatively, the patient’s eGFR met criteria for a diagnosis of CKD prior to the AKI alert.
Cause of AKI	The clinician documented an AKI cause in a visit note in the EHR within two weeks of the AKI alert.
Hypovolemia	The clinician noted hypovolemia, volume depletion, overdiuresis, diarrhea or vomiting as the cause of AKI.
Cardiorenal	The clinician noted cardiorenal syndrome as the cause of AKI.
Drugs (including contrast)	The clinician noted drugs, contrast, tacrolimus toxicity, bactrim toxicity, levaquin toxicity or nephrotoxins as the cause of AKI. Diuretics were not considered nephrotoxins. Instead diuretics or overdiuresis was categorized as hypovolemia.
Other	The clinician noted obstructive uropathy, progression to CKD, sepsis, rhabdomyolysis, upper GI bleeding, membranous nephropathy, hepatorenal syndrome, urinary tract infection, radical prostatectomy, hypotension or hypercalcemia as the cause of AKI.
Multifactorial	There was more than one documented cause of AKI.

*If the definition is met, the corresponding field in the standardized review instrument is marked as “1”. If the definition is not met, the field is marked as “0”.

AKI = acute kidney injury; EHR = electronic health record; CKD = chronic kidney disease

Supplemental Table of Contents

Supplemental Table 1. Definitions of Parameters Collected During Medical Record Review

Supplemental Table 1. Definitions of Parameters Collected During Medical Record Review

Parameter	Definition*
Documented AKI	Any clinician acknowledged AKI in the EHR in a visit note (any visit note, not limited to nephrology) or patient communication within two weeks of the AKI alert.
Documented Clinical Action	Any clinician documented one or more of the clinical actions below in the EHR in a visit note (any visit note, not limited to nephrology) or patient communication within two weeks of the AKI alert.
Repeat Labs	Any clinician noted that laboratory testing should be repeated.
Encourage Hydration	Any clinician noted that the patient should hydrate, increase fluid intake or present to a care facility for intravenous hydration.
Discontinue Medications	Any clinician noted that the patient should stop taking a medication.
Change Dosing	Any clinician noted that the patient should change the dosing of a medication or stop taking a medication temporarily. The temporary hold was included in “change dosing” because the authors intended for the “discontinue medications” field to reflect permanent or indefinite discontinuation of a medication.
Nephrology Consult	A nephrology consult occurred (as evidenced by a nephrology visit note) within 2 weeks of the AKI alert. Alternatively, there was documentation that the patient had a nephrology consult at an outside facility within 2 weeks of the AKI alert.
Hospitalization	The patient was hospitalized specifically for evaluation and management of AKI. The following criteria had to be met: 1) patient was admitted to observation or inpatient status at a Mass General Brigham inpatient facility and 2) the admission occurred after the AKI alert. At least one instance of documentation from that admission, e.g. emergency department expect note or inpatient history and physical, listed AKI as the primary problem, chief complaint, or reason for hospitalization.
Underlying CKD	The clinician noted in the EHR that the patient had a diagnosis of CKD in a visit note, patient communication, or problem list prior to the AKI alert. Alternatively, the patient’s eGFR met criteria for a diagnosis of CKD prior to the AKI alert.
Cause of AKI	The clinician documented an AKI cause in a visit note in the EHR within two weeks of the AKI alert.
Hypovolemia	The clinician noted hypovolemia, volume depletion, overdiuresis, diarrhea or vomiting as the cause of AKI.
Cardiorenal	The clinician noted cardiorenal syndrome as the cause of AKI.
Drugs (including contrast)	The clinician noted drugs, contrast, tacrolimus toxicity, bactrim toxicity, levaquin toxicity or nephrotoxins as the cause of AKI. Diuretics were not considered nephrotoxins. Instead diuretics or overdiuresis was categorized as hypovolemia.
Other	The clinician noted obstructive uropathy, progression to CKD, sepsis, rhabdomyolysis, upper GI bleeding, membranous nephropathy, hepatorenal syndrome, urinary tract infection, radical prostatectomy, hypotension or hypercalcemia as the cause of AKI.
Multifactorial	There was more than one documented cause of AKI.

*If the definition is met, the corresponding field in the standardized review instrument is marked as “1”. If the definition is not met, the field is marked as “0”.

AKI = acute kidney injury; EHR = electronic health record; CKD = chronic kidney disease